The history of innovation in Radio Astronomy

From Antikythera to the SKA Kerastari, Greece 12 Jun 2012



Ron Ekers CSIRO, Australia

## Galileo Galilei - 1609

• Galileo builds a telescope and he sees the moons of Jupiter.



*"Four planets, never seen since the beginning of the World right up to our day"* 



#### Karl Jansky Bell Telephone Laboratory 1932



## Beginning of Radio Astronomy

- Jansky's Cosmic Hiss
  - Unexpected source of noise peaking each day
  - signal arrives 4 min earlier each day
  - reaction from Bell Labs "so faint not even interesting as a source of radio interference!"
  - not accepted by the astronomical community at the time
  - Jansky died in 1944 before the importance
    - of his discovery was appreciated



- Dec 1938
- Edmond Hamilton
  - The Cosmic Hiss
  - Based on Jansky's observation of cosmic hiss
- Extra Terrestrial Intelligence?

#### Science Fiction





1 May 2005



Grote Reber 1911-2002 Wheaton Illinois - 1937 - 32' parabolic dish Cosmic radiation had to be nonthermal – No theory until 1950



# The Dish

#### Parkes 64m, Australia 1960

ers - APRIM2011

#### What are the Radio Stars? Cliff Interferometer - 1948 • Bolton, Stanley and Slee (CSIRO, Australia) – 100MHz Yagi



3 Mar 200520June2007

WW2 radar used interference from reflected signals to measure elevation



What are the Radio Stars? Cliff Interferometer - 1948 • Bolton, Stanley and Slee (CSIRO, Australia) – 100MHz Yagi



Used interference fringes to measure source position



R D EkersR. Ekers

3 Mar 200520June2007

S I

RO





#### Centaurus A ATCA Mosaic





#### Centaurus A ATCA Mosaic

#### Today we use Aperture Synthesis **Images routinely**

#### But the path to developing the underlying concepts has a rich history involving discovery, sociology, and some incredible. individuals Radio Image of

Ionised Hydrogen in Cyg X CGPS (Penticton)

## History of Fourier Synthesis Imaging

- 1891: Michelson defines fringe visibility
  - Gives the Fourier equations but doesn't call it a Fourier transform
- Stereo X-ray imaging
- 1912: X-ray diffraction in crystals
- 1930: van Cittert-Zernike theorem
  - Now considered the basis of Fourier synthesis imaging
  - Played no role in the early radio astronomy developments but appears in the literature after Born & Wolf *Principles* of Optics (1960)
- 1930-38: 3D X-ray tomography
  - Analogue devices to do back projection summation

## X-ray Crystallography

- 1912
  - X-ray diffraction in crystals
- 1936
  - Lipson & Beevers strips
  - Fourier synthesis calculations routine in X-ray crystallography



• 1939

SIRO

- Bragg's X-ray crystallography group flourishing at the Cavendish Laboratory
  - 2D Fourier analysis
  - phase problem,



16

# $Ratcliffe \Rightarrow Pawsey \Rightarrow Bracewell$ $Cambridge \Rightarrow Sydney$

#### • 1935

- Pawsey PhD with Ratcliffe at Cambridge (ionosphere)

• 1940

 Pawsey joins CSIRO Radiophysics Laboratory in Sydney but maintains strong links with Ratcliffe in Cambridge

#### • 1946-1949

Pawsey introduces Bracewell to duality of physical and mathematical descriptions following Ratcliffe's style
Bracewell sent from Sydney to work with Ratcliffe



## Ryle and the Cavendish

- 1945
  - Ryle joins Cavendish laboratory
    - uses WWII radar technology for radio astronomy
- 1946
  - Ryle and Vonberg (Nature 158, 339-340 (Aug 1946)
    - interferometric measurement of sunspots



 introduces the use of a Michelson interferometer to measure the angular diameter of the source of the radiation and references Michelson

#### McCready, Pawsey & Payne-Scott 1947

- Proc Roy Soc, Aug 1947 received July 1946!
- Used the phase of the sea interferometer fringes (lobes) to co-locate solar emission with sunspots
- They note that its possible in principal to determine the actual distribution by Fourier synthesis using the phase and amplitude at a range of height or wavelength.
- They consider using wavelength as a suitable variable as unwise since the solar bursts are likely to have frequency dependent structure.



They note that getting a range of cliff height is clumsy and suggest a different interference method would be more practical.

#### Fourier Transforms - 1953

- Lipson-Beevers strips
  - 25x25 array to 2 digits 1 person in 24 hours
- Punched card tabulator
  - 25x25 array to 3 digits in 8 hours (4 operators!)





M. Ryle and Ann C. Neville



## First Cambridge Earth Rotation Synthesis Image

- Ryle & Neville, MNRAS 1962
- June 1961
- North pole survey
- 4C aerials
- 178 MHz
- Not superseded until LOFAR & MWA!

#### Cambridge One-Mile Telescope:







## Nobel Prize 1974 Sir Martin Ryle

from the presentation "The radio-astronomical instruments invented and developed by Martin Ryle, and utilized so successfully by him and his collaborators in their observations, have been one of the most important elements of the latest discoveries in Astrophysics."



### Westerbork: 1970

- Hogbom (Cambridge)
- Christiansen (Sydney)

#### Benelux cross $\rightarrow$ WSRT

- 12 x 25m dishes 1.5km
  - Two moveable
  - 10 redundant spacings
  - Self calibration
  - Two more dishes at 3km added later



# VLA

# New Mexico

#### Self Calibration

- 1958: Phase and amplitude closure
  - Jennison (Jodrell Bank)
- 1977: Redundant spacing interferometry
  - Hamaker, O'Sullivan, Noordam (Westerbork)
- 1974-79: Phase closure in VLBI imaging
  - Rogers, Yee, Readhead, Cotton....
- 1980: Antenna based calibration
  - Clark, Schwab (VLA)
- 1983: Self cal ≡ phase closure ≡ adaptive optics
   Cornwell

#### Little Science – Big Science

Derek J. de Solla Price (1963) *Little Science, Big Science* Columbia University Press





27 Nov 1999

R D Ekers - APRIM2011

#### Derek J de Solla Price

- Price worked as a teacher of applied mathematics at Raffles College ( University of Singapore) in 1948.
- It was there that he formulated his theory on the exponential growth of science
- The idea occurred to him when he noticed the growth in stacks of the *Philosophical Transactions of the Royal Society* between 1665 and 1850
- He had the complete set in his home while Raffles College had its library built.



## Radio Telescope Sensitivity

- Exponential increase in sensitivity x  $10^5$ since 1940 !
  - 3 year doubling time for sensitivity



27 Nov 1999

SIRO

#### The Original Moore's Law Plot

 In 1965 Gordon Moore (co-founder of Intel) noted that the transistor density of semiconductor chips doubled roughly every 18 months.

27 Nov 1999





Electronics, April 1965

### Microprocessor performance

 Moore's Law
 – Intel 2000





R D Ekers - APRIM2011

- De Solla Price (1963): most scientific advances follow laboratory experiments
  - exponential growth in capability occurs in all successful science
  - Little science, big Science
     Individual researcher





- De Solla Price (1963): most scientific advances follow laboratory experiments
  - exponential growth in capability occurs in all successful science
  - Little science, big Science
     Individual researcher
     Institute laboratory





- De Solla Price (1963): most scientific advances follow laboratory experiments
  - exponential growth in capability occurs in all successful science
  - Little science, big Science
     Individual researcher
     Institute laboratory
     National Facility





- De Solla Price (1963): most scientific advances follow laboratory experiments
  - exponential growth in capability occurs in all successful science
  - Little science, big Science
     Individual researcher
    - ↓ Institute laboratory ↓ National Facility



RO



# SKA 2020?